

NEWS RELEASE

1111 North Fairfax Street
Alexandria, VA 22314-1488
703 684 2782
703 684 7343 fax
www.apta.org

CONTACT:

Jennifer Rondon
Public and Media Relations
American Physical Therapy Association
703/706-3216

STUDY FINDS PHYSICAL THERAPY JUST AS EFFECTIVE AS SURGERY IN PATIENTS WITH A TORN MENISCUS AND ARTHRITIS OF THE KNEE

Results Should Help Change Practice, Says APTA

Alexandria, VA, March 21, 2013 – A *New England Journal of Medicine* (NEJM) study showing that physical therapy is just as effective as surgery in patients with meniscal tears and arthritis of the knee should encourage many health care providers to reconsider their practices in the management of this common injury, according to the American Physical Therapy Association (APTA).

The study, published March 19, showed no significant differences in functional improvement after 6 months between patients who underwent surgery with postoperative physical therapy and those who received standardized physical therapy alone.

“This study demonstrates what physical therapists have long known,” explained APTA President Paul A. Rockar Jr, PT, DPT, MS. “Surgery may not always be the best first course of action. A physical therapist, in many cases, can help patients avoid the often unnecessary risks and expenses of surgery. This study should help change practice in the management of symptomatic meniscal tears in patients with knee osteoarthritis.”

According to lead physical therapist for the trial and American Physical Therapy Association (APTA) member Clare Safran-Norton, PT, PhD, OCS, “our findings suggest that a course of physical therapy in this patient population may be a good first choice since there were no group differences at 6 months and 12 months in this trial. These findings should help surgeons, physicians, physical therapists, and patients in decision-making regarding their treatment options.”

Researchers at 7 major universities and orthopedic surgery centers around the country studied 351 patients aged 45 years or older who had a meniscal tear and mild-to-moderate osteoarthritis of the knee. Patients were randomly assigned to groups who received either surgery and postoperative physical therapy or standardized physical therapy. Within 6-12 months, patients who had physical therapy alone showed similar improvement in functional status and pain as those who had undergone arthroscopic partial meniscectomy surgery.

—more—

Patients who were given standardized physical therapy—individualized treatment and a progressive home exercise program—had the option of “crossing over” to surgery if substantial improvements were not achieved. Thirty percent of patients crossed over to surgery during the first 6 months. At 12 months these patients reported similar outcomes as those who initially had surgery. Seventy percent of patients remained with standardized physical therapy.

According to an accompanying editorial in *NEJM*, “millions of people are being exposed to potential risks associated with a treatment [surgery] that may or may not offer specific benefit, and the costs are substantial.” Physical therapist and APTA member Mary Ann Wilmarth, PT, DPT, MS, OCS, MTC, Cert MDT, chief of physical therapy at Harvard University, said, “Physical therapists are experts in improving mobility and restoring motion. The individualized treatment approach is very important in the early phases of rehabilitation in order to achieve desired functional outcomes and avoid setbacks or complications.”

The American Physical Therapy Association (APTA) represents more than 85,000 physical therapists, physical therapist assistants, and students of physical therapy nationwide. Learn more about meniscal tears and other conditions that physical therapists can treat, and find a physical therapist in your area at www.MoveForwardPT.com. Consumers are encouraged to follow us on Twitter ([@MoveForwardPT](https://twitter.com/MoveForwardPT)) and [Facebook](https://www.facebook.com/MoveForwardPT).

###